

(FILE 'HOME' ENTERED AT 12:00:51 ON 08 NOV 2007)

FILE 'BIOSIS, SCISEARCH, AGRICOLA, MEDLINE' ENTERED AT 12:01:19 ON 08 NOV 2007

L1 15614 S AZUMA, ?/AU  
L2 267 S HAMAMATSU, ?/AU  
L3 2631 S FUJINAGA, ?/AU  
L4 836 S (CELL WALL SKELETON)  
L5 49989 S (BCG OR BACILLUS CALMETTE GUERIN)  
L6 197 S L1 AND L4  
L7 0 S L2 AND L4  
L8 0 S L3 AND L4  
L9 323 S L4 AND L5  
L10 461011 S (EXTRACTION)  
L11 2 S L10 AND L9  
L12 79952 S (EMULSION)  
L13 2417 S L12 AND L10  
L14 1 S L13 AND L9  
L15 370065 S (SOLVENT)  
L16 47525 S L10 AND L15  
L17 2 S L16 AND L4

=>

**Brief Communication**

**Tumor Suppression by Cell Walls of  
*Mycobacterium bovis* Attached to Oil  
Droplets<sup>1</sup>**

Berton Zbar, Herbert J. Rapp, and Edgar E. Ribi,<sup>2</sup> *Biology  
Branch, National Cancer Institute,<sup>3</sup> Bethesda, Maryland 20014*

**SUMMARY**—The growth of syngeneic guinea-pig tumor transplants in skin was suppressed if the tumor cells were inoculated together with bacillus Calmette-Guérin (BCG) cell walls attached to oil droplets. Tumor growth was not inhibited if the tumor cells were given together with 1) BCG cell walls, 2) oil droplets, and 3) oil droplets and BCG cell walls prepared so as to prevent attachment of the walls to the oil droplets. Animals in which tumor growth was suppressed acquired systemic tumor immunity.—*J Nat Cancer Inst* 48: 831-835, 1972.

WE HAVE reported that the intradermal growth of a transplantable, syngeneic guinea-pig hepatocarcinoma in normal guinea pigs is suppressed if the tumor cells are inoculated together with living bacillus Calmette-Guérin (BCG) (1, 2). Heat-killed BCG or extracts of tubercle bacilli did not inhibit tumor growth. For treatment of cancer patients, however, nonliving mycobacterial preparations with tumor-suppressive properties would be preferable to living BCG. We now report that cell walls of BCG attached to oil droplets possess potent tumor-suppressive activity. Preparations of this type are as effective as living BCG in preventing pulmonary tuberculosis in mice and monkeys (3, 4).

**MATERIALS AND METHODS**

**Animals.**—Sewall-Wright inbred strain-2 male guinea pigs were obtained from the Laboratory Aids Branch, Division of Research Services, National Institutes of Health (5). Skin grafts between

members of the strain are not rejected (6). Guinea pigs were grouped 6 per cage and fed Wayne guinea pig chow daily and kale 3 times a week.

**Tumors.**—We have described the induction of primary hepatomas in strain-2 guinea pigs fed the water-soluble carcinogen diethylnitrosamine (7) and the antigenic and biologic properties of the transplantable tumors derived from the primary hepatomas (8, 9). A transplantable hepatoma designated tumor line 10 (seventh to eleventh transplant generations) was used. Tumor line 10 is a hepatocellular carcinoma converted to ascites form. Intradermal inoculation of  $10^5$  ascites line-10 tumor cells leads to progressive intradermal growth and metastases to the regional lymph nodes. Animals

<sup>1</sup>Received November 16, 1971; accepted January 3, 1972.

<sup>2</sup>Rocky Mountain Laboratory, National Institutes of Allergy and Infectious Disease, Hamilton, Mont.

<sup>3</sup>National Institutes of Health, Public Health Service, U.S. Department of Health, Education, and Welfare.

die about 60 days after intradermal inoculation of  $1.5 \times 10^6$  tumor cells (2).

**Preparation of BCG cell walls.**—Cell walls of the Glaxo strain of BCG were prepared as in (10). Mycobacteria were grown in Sauton's medium at the Lederle Laboratories (Pearl River, N.Y.). After 3 weeks in culture, the bacteria were disrupted in the Sorvall refrigerated pressure cell at 35,000 pounds per square inch. Cell walls were obtained by centrifugation of the mixture at  $20,000 \times g$  for 1 hour. The pellet was resuspended in distilled water and washed several times until only cell walls were visible under the electron microscope.

**Preparation of BCG cell walls attached to oil droplets.**—Freeze-dried cell walls (25 mg) were placed in a 15-ml tissue grinder equipped with a Teflon pestle (Scientific Glass Apparatus Co., Inc., Bloomfield, N.J.). A light mineral oil (0.12 ml), Drakeol 6 VR (Pennsylvania Refining Co., Butler, Penna.), was added to the cell walls. This mixture was ground to a smooth paste; the pestle was rotated at 800 rpm. Ten ml 0.85% NaCl containing 0.2% Tween 80 (polyoxyethylene derivative of sorbitan mono-oleate, Atlas Powder Co., Wilmington, Del.) was added to the paste, and grinding was continued until a well-dispersed oil-in-water emulsion was obtained. The emulsion was poured into a sterile container and 6.7 ml Tween-saline diluent was added to the grinding tube. Grinding was continued for 2–3 minutes. The 2 emulsions were combined and ground for an additional 2–3 minutes. This emulsion containing 1.5 mg cell walls per ml was heated at  $65^\circ\text{C}$  for 30 minutes in a water bath. Examination of the product with the light microscope showed spherical oil droplets varying from  $<1 \mu$  to  $>15 \mu$ ; discrete particles appeared to be attached to the surface of the oil droplets. Photographs of similar emulsions are in (11).

**Preparation of oil droplets alone, cell walls alone, and cell walls not attached to oil droplets.**—Oil droplets alone: This preparation was made in the same way as the BCG cell walls attached to oil droplets except for the omission of the cell walls. Under the light microscope this product showed clear spherical oil droplets varying from  $<1 \mu$  to  $>15 \mu$ . Cell walls alone: The method was the same as for the BCG cell walls attached to oil droplets except for the omission of the oil. Under the light microscope

this product showed cell wall fragments varying from  $2\text{--}5 \mu$ . Cell walls not attached to oil droplets: In this preparation, the oil in water emulsion was made first and the cell walls were added without further grinding. The cell walls were dispersed in the emulsion by gentle mixing. Under the light microscope this product showed clear spherical oil droplets of varying size and aggregated cell walls that remained unattached to oil droplets.

**Preparation of tumor cell mixtures for inoculation.**—Ascites tumor cells, obtained as described in (12), were washed 3 times in medium 199 without antibiotics. Equal volumes of ascites tumor cells and the appropriate emulsion or cell wall preparation were mixed at room temperature in  $13 \times 100$  mm siliconized, sterile, glass test tubes; they were incubated at  $37^\circ\text{C}$  for 10 minutes, and then 0.1 ml was injected intradermally into unimmunized guinea pigs.

**Measurement of delayed skin reactions and papules.**—Delayed skin reactions were measured 24 hours after injection of the mixtures. Papules were measured at several intervals after injection. Two diameters of the skin reaction or papule perpendicular to one another were measured. The two diameters were used to calculate the square of the average radius of the skin reaction or papule. In experiments in which several animals were challenged with portions of the same tumor-cell mycobacterial antigen mixture, the square of the average radius for each lesion produced by the same number of tumor cells was calculated. Results were expressed as the mean of the square  $\pm$  the standard error of the mean.

Each experimental group contained 3 guinea pigs. At each time point, the results represent the mean value of 3 animals. Each animal received a single test mixture.

## RESULTS

In preliminary experiments, oil-treated cell walls of BCG were injected intradermally into guinea pigs immunized with BCG and into unimmunized guinea pigs. The following day, typical delayed cutaneous-hypersensitivity reactions were seen only in guinea pigs immunized to BCG. Three days later, in immunized and unimmunized guinea pigs an inflammatory nodule was at the injection site of

oil-treated  
for 3 w  
healed  
oil-treated  
bacteria  
chronic  
weeks.

The  
treated  
experi  
injection  
and 1.  
was gi  
droplet  
line-10  
derma  
10 cc  
0.2%  
with 1  
with (1  
the c  
Oil-tr  
growth  
contain  
The  
deterr  
droplet  
the ce  
for tu

PAPULE SIZE, RADIUS SQUARED, mm<sup>2</sup>

Tumor  
treated  
drop  
+ 1  
taint  
Δ:

oil-treated cell walls of BCG. This nodule persisted for 3 weeks. A small scar was left at the site of the healed nodule. This experiment indicated that oil-treated cell walls of BCG contained mycobacterial antigens and, like living BCG, produced a chronic inflammatory nodule which healed in 3-4 weeks.

The possible tumor-suppressive properties of oil-treated cell walls were evaluated in the next experiment. Group 1 received an intradermal injection containing oil-treated cell walls (75  $\mu$ g) and  $1.5 \times 10^6$  living line-10 tumor cells. Group 2 was given an intradermal injection containing oil droplets without cell walls and  $1.5 \times 10^6$  living line-10 tumor cells. Group 3 received an intradermal injection containing  $1.5 \times 10^6$  living line-10 cells diluted with 0.85% saline containing 0.2% Tween. Group 4 was inoculated intradermally with  $1.5 \times 10^6$  living line-10 tumor cells diluted with 0.85% saline without Tween. The results of the experiment are illustrated in text-figure 1. Oil-treated cell walls completely suppressed tumor growth. Oil droplets without cell walls or saline containing Tween did not inhibit tumor growth.


The following experiment was designed to determine whether bacterial cell walls without oil droplets would inhibit tumor growth and whether the cell walls had to be attached to the oil droplets for tumors to be suppressed. Group 1 received an

intradermal injection of 75  $\mu$ g oil-treated cell walls +  $1.5 \times 10^6$  living line-10 tumor cells. Group 2 was given 75  $\mu$ g cell walls without oil droplets +  $1.5 \times 10^6$  living line-10 tumor cells. Group 3 received 75  $\mu$ g cell walls not attached to oil droplets +  $1.5 \times 10^6$  line-10 cells. Group 4 was treated with  $1.5 \times 10^6$  line-10 cells diluted with saline containing Tween. The results of this experiment (text-fig. 2) indicate that cell walls alone and cell walls in the presence of, but not attached to, oil droplets were ineffective in tumor inhibition.


To test whether guinea pigs treated with a mixture of oil-treated cell walls of BCG and tumor cells had developed systemic tumor immunity, guinea pigs received intradermal injections of  $10^6$  living line-10 tumor cells. The challenge dose was inoculated contralateral to the immunization site. The presence of delayed cutaneous hypersensitivity reactions to tumor cells and prevention of tumor growth (table 1) indicate that guinea pigs treated with vaccine had developed systemic tumor immunity.

## DISCUSSION

These experiments demonstrate that a nonliving mycobacterial preparation can completely suppress tumor growth. The tumor-suppressive property of oil-treated cell walls is in contrast to the previously



TEXT-FIGURE 1.—Suppression of tumor growth by oil-treated cell wall of BCG. ●: Cell walls attached to oil droplets +  $1.5 \times 10^6$  line-10 tumor cells. ○: Oil droplets +  $1.5 \times 10^6$  line-10 tumor cells. □: 0.85% saline containing 0.2% Tween 80 +  $1.5 \times 10^6$  line-10 tumor cells. Δ: 0.85% saline +  $1.5 \times 10^6$  line-10 tumor cells.



TEXT-FIGURE 2.—Suppression of tumor growth by oil-treated cell walls of BCG. ●: Cell walls attached to oil droplets +  $1.5 \times 10^6$  line-10 tumor cells. ○: Cell walls +  $1.5 \times 10^6$  line-10 tumor cells. □: 0.85% saline containing 0.2% Tween 80 +  $1.5 \times 10^6$  line-10 tumor cells. Δ: Cell walls not attached to oil droplets +  $1.5 \times 10^6$  line-10 tumor cells.

TABLE 1.—Tumor immunity after intradermal injection of oil-treated mycobacterial cell walls and living tumor cells

| Treatment                                     | Delayed cutaneous hypersensitivity reactions at 24 hr—<br>radius squared, mm <sup>2</sup> |          | Tumor incidence at 10 days<br>after challenge with 10 <sup>6</sup> tumor<br>cells |         |
|---|---|----------|---|---------|
|   | Expt. 1*  | Expt. 2  | Expt. 1   | Expt. 2 |
| Oil-treated cell walls and living tumor cells | 27 ± 2  | Not done | 0/3   | 0/3     |
| None  | 7 ± 1   | Not done | 6/6   | 5/5     |

\*In expt. 1, guinea pigs were challenged the 28th day after immunization and in expt. 2 the 22d day after immunization. Immunized animals were free of tumor at the time of this report, 2 months after challenge. Tumors in the unimmunized group grew progressively.

reported (2, 13) lack of tumor inhibition in guinea pigs not immunized to BCG when other nonliving mycobacterial antigen preparations were used (cell walls alone, purified protein derivative, heat-killed BCG, crude tuberculo-protein). In immunized guinea pigs, tumor may be partially suppressed by these nonliving antigens. There have been reports that nonliving preparations or fractions of mycobacteria are effective in tumor inhibition (14). The experimental model described in this report offers the possibility of comparing the tumor-inhibiting capacity of nonliving preparations of mycobacteria.

Injection of oil-treated cell walls mixed with living tumor cells leads to the development of tumor immunity. Recent studies of tumor immunity produced by injection of mycobacterial antigen-tumor cell mixtures indicate that differences exist in the intensity of tumor immunity produced by different mycobacterial antigen preparations (15). For example, intradermal immunization with complete Freund's adjuvant and tumor cells was not as effective as intradermal immunization with living BCG and tumor cells.

We do not know why BCG cell walls attached to oil droplets can suppress tumor growth. If the mechanism of tumor suppression by oil-treated BCG cell walls is similar to that obtained with living BCG, this means that in some unknown fashion the oil droplets make it possible for the recipient to respond immunologically to BCG cell-wall antigens by the production of specifically sensitized lymphocytes. Interaction of these lymphocytes and BCG antigens in the presence of tumor cells causes the accumulation of nonlymphoid cells responsible for tumor cell death.

This work may provide a simple, rapid model for determining the components of the tubercle bacillus required for tumor suppression and indi-

cates that a nonliving mycobacterial antigen preparation may be clinically useful for inhibiting tumor growth.

#### REFERENCES

- (1) ZBAR B, BERNSTEIN I, TANAKA T, et al: Tumor immunity produced by the intradermal inoculation of living tumor cells and living *Mycobacterium bovis* (strain BCG) *Science* 170:1217-1218, 1970
- (2) ZBAR B, BERNSTEIN ID, RAPP HJ: Suppression of tumor growth at the site of infection with living bacillus Calmette-Guérin. *J Nat Cancer Inst* 46:831-839, 1971
- (3) RIBI E, LARSON C, WIGHT W, et al: Effective non-living vaccine against experimental tuberculosis in mice. *J Bact* 91:975-983, 1966
- (4) RIBI E, ANACKER RL, BARCLAY WR, et al: Efficacy of mycobacterial cell walls as a vaccine against airborne tuberculosis in the rhesus monkey. *J Infect Dis* 123:527-538, 1971
- (5) LOEB L, WRIGHT S: Transplantation and individuality differentials in inbred families of guinea-pigs. *Amer J Path* 3:251-283, 1927
- (6) BAUER JA Jr: Histocompatibility in inbred strains of guinea pigs. *Ann NY Acad Sci* 73:663-672, 1958
- (7) RAPP HJ, CHURCHILL WH Jr, KRONMAN BS, et al: Antigenicity of a new diethylnitrosamine-induced transplantable guinea pig hepatoma: Pathology and formation of ascites variant. *J Nat Cancer Inst* 41:1-11, 1968
- (8) ZBAR B, WEPF HT, RAPP HJ, et al: Antigenic specificity of hepatomas induced in strain-2 guinea pigs by diethylnitrosamine. *J Nat Cancer Inst* 43:833-841, 1969
- (9) ZBAR B, WEPF HT, RAPP HJ, et al: Transplantable hepatomas induced in strain-2 guinea pigs by diethylnitrosamine: Characterization by histology, growth, and chromosomes. *J Nat Cancer Inst* 43:821-831, 1969
- (10) RIBI E, ANACKER RL, BREHMER W, et al: Factors influencing protection against experimental tuberculosis in mice by heat-stable cell wall vaccines. *J Bact* 92:869-879, 1966

- (11) BARCLAY WR, ANACKER R, BREZMER W, et al: Effects of oil-treated mycobacterial cell walls on the organs of mice. *J Bact* 94:1736-1745, 1967
- (12) CHURCHILL WH JR, RAPP HJ, KRONMAN BS, et al: Detection of antigens of a new diethylnitrosamine-induced transplantable hepatoma by delayed hypersensitivity. *J Nat Cancer Inst* 41:19-29, 1968
- (13) ZBAR B, WEPSIC HT, BORROS T, et al: Tumor-graft rejection in syngeneic guinea pigs: Evidence for a two-step mechanism. *J Nat Cancer Inst* 44:473-481, 1970
- (14) WEISS DW, BONEAC NS, LESLIE P: Studies on the heterologous immunogenicity of a methanol-insoluble fraction of attenuated tubercle bacilli (BCG). II. Protection against tumor isografts. *J Exp Med* 124:1039-1065, 1966
- (15) BARTLETT G, ZBAR B: Unpublished observations